

May 2023



COMMENTARY BY THE HEI REVIEW COMMITTEE SUMMARIZING AND EVALUATING THE INVESTIGATORS' REPORT

Ambient Air Pollution and All-Cause and Cause-Specific Mortality in an Analysis of Asian Cohorts

Downward and Vermuelen

Health Effects Institute

Ambient Air Pollution and All-Cause and Cause-Specific Mortality in an Analysis of Asian Cohorts

George S. Downward and Roel Vermeulen

with a Commentary by HEI's Review Committee

Research Report 213 Health Effects Institute Boston, Massachusetts

Trusted Science · Cleaner Air · Better Health

CONTENTS

About HEI	iii				
Contributors					
COMMENTARY by the Review Committee	1				
INTRODUCTION	1				
Scientific background	2				
Study Population Exposure Assessment Outcome Assessment Analyses Summary of Results Across Cohorts Summary of Results Within Cohorts	2 2 2 2 4 4				
HEI REVIEW COMMITTEE'S EVALUATION	4				
Inadequate Adjustment for Characteristics That Likely Correlate with Air Pollution and Mortality Heterogeneity in Effect Estimates Substantial Temporal and Spatial Misalignment of the Exposure Data Household Air Pollution Was Not Examined Broader Context of Air Pollution and Health In Asia	6 7 8 8				
	7				
REFERENCES	10				
Abbreviations and Other Terms	13				
HEI Board, Committees, and Staff	14				

ABOUT HEI

The Health Effects Institute is a nonprofit corporation chartered in 1980 as an independent research organization to provide high-quality, impartial, and relevant science on the effects of air pollution on health. To accomplish its mission, the Institute

- · Identifies the highest-priority areas for health effects research
- Competitively funds and oversees research projects
- · Provides intensive independent review of HEI-supported studies and related research
- Integrates HEI's research results with those of other institutions into broader evaluations
- Communicates the results of HEI's research and analyses to public and private decision makers.

HEI typically receives balanced funding from the U.S. Environmental Protection Agency and the worldwide motor vehicle industry. Frequently, other public and private organizations in the United States and around the world also support major projects or research programs. HEI has funded more than 340 research projects in North America, Europe, Asia, and Latin America, the results of which have informed decisions regarding carbon monoxide, air toxics, nitrogen oxides, diesel exhaust, ozone, particulate matter, and other pollutants. These results have appeared in more than 260 comprehensive reports published by HEI, as well as in more than 2,500 articles in the peer-reviewed literature.

HEI's independent Board of Directors consists of leaders in science and policy who are committed to fostering the public–private partnership that is central to the organization. The Research Committee solicits input from HEI sponsors and other stakeholders and works with scientific staff to develop a Five-Year Strategic Plan, select research projects for funding, and oversee their conduct. The Review Committee, which has no role in selecting or overseeing studies, works with staff to evaluate and interpret the results of funded studies and related research.

All project results and accompanying comments by the Review Committee are widely disseminated through HEI's website (*www.healtheffects.org*), reports, newsletters and other publications, annual conferences, and presentations to legislative bodies and public agencies.

CONTRIBUTORS

RESEARCH COMMITTEE

David A. Savitz, Chair Professor of Epidemiology, School of Public Health, and Professor of Obstetrics and Gynecology, Alpert Medical School, Brown University

David C. Dorman Professor, Department of Molecular Biomedical Sciences, College of Veterinary Medicine, North Carolina State University

Christina H. Fuller Associate Professor, School of Environmental, Civil, Agricultural and Mechanical Engineering, University of Georgia College of Engineering

Marianne Hatzopoulou Professor, Civil and Mineral Engineering, University of Toronto, Canada, Research Chair in Transport Decarbonization and Air Quality

Amy H. Herring Sara & Charles Ayres Professor of Statistical Science and Global Health, Duke University

Heather A. Holmes Associate Professor, Department of Chemical Engineering, University of Utah

REVIEW COMMITTEE

Melissa J. Perry, Chair Dean, College of Health and Human Services, George Mason University

Sara D. Adar Associate Professor and Associate Chair, Department of Epidemiology, University of Michigan School of Public Health

Kiros T. Berhane Professor and Chair, Department of Biostatistics, Mailman School of Public Health, Columbia University

Ulrike Gehring Associate Professor, Institute for Risk Assessment Sciences, Utrecht University

Michael Jerrett Professor and Chair, Department of Environmental Health Sciences, Fielding School of Public Health, University of California–Los Angeles

Frank Kelly Henry Battcock Chair of Environment and Health and Director of the Environmental Research Group, Imperial College London School of Public Health, United Kingdom

HEI PROJECT STAFF

Pallavi Pant Head of Global Health (Study Oversight)

Eleanne van Vliet Staff Scientist (Study Oversight)

Hanna Boogaard Consulting Principal Scientist (Report Review and Commentary)

Yi Lu Staff Scientist (Report Review)

Neil Pearce Professor of Epidemiology and Biostatistics, London School of Hygiene and Tropical Medicine, United Kingdom

Ana M. Rule Assistant Professor and Director, Environmental Exposure Assessment Laboratories, Department of Environmental Health and Engineering, Johns Hopkins School of Public Health

Evangelia (Evi) Samoli Associate Professor of Epidemiology and Medical Statistics, Department of Hygiene, Epidemiology and Medical Statistics, School of Medicine, National and Kapodistrian University of Athens, Greece

Neeta Thakur Associate Professor of Medicine, University of California, San Francisco

Gregory Wellenius Professor, Department of Environmental Health, Boston University School of Public Health

Jana B. Milford Professor Emerita, Department of Mechanical Engineering and Environmental Engineering Program, University of Colorado–Boulder

Jennifer L. Peel Professor of Epidemiology, Colorado School of Public Health and Department of Environmental and Radiological Health Sciences, Colorado State University

Eric J. Tchetgen Tchetgen Luddy Family President's Distinguished Professor, Professor of Statistics and Data Science, The Wharton School, University of Pennsylvania

John Volckens Professor, Department of Mechanical Engineering, Walter Scott Jr. College of Engineering, Colorado State University

Kristin EcklesSenior Editorial ManagerHope GreenEditorial Project ManagerMary BrennanConsulting Editor

COMMENTARY Review Committee

HE

Research Report 213, *Ambient Air Pollution and All-Cause and Cause-Specific Mortality in an Analysis of Asian Cohorts*, G. S. Downward and R. Vermeulen

INTRODUCTION

Air pollution is a major global public health risk factor. There is now broad expert consensus that exposure to air pollution causes an array of adverse health effects based on evidence from a large body of scientific literature that has grown exponentially since the mid-1990s (IARC 2016; Thurston et al. 2017; U.S. EPA 2016, 2019; WHO 2021).

Based on that evidence, the Global Burden of Disease (GBD*) project estimated that in 2019 air pollution ranked as the leading environmental risk factor for global mortality, surpassed only by high blood pressure, tobacco use, and poor diet (HEI 2020). The air pollution burden varies widely around the globe, and is highest in countries in Asia and Africa, partly due to the typically high exposure levels in those regions.

Much of what is currently known about the adverse effects of ambient air pollution comes from studies conducted in high-income regions, especially North America and Europe, with relatively low air pollution levels. Studies of long-term exposure and morbidity and mortality in low- and middle-income countries have emerged more recently. Hence, an integrated exposure-response (IER) function was developed to estimate mortality relative risks across the global exposure range and has been used by the GBD collaboration and the World Health Organization (WHO) to estimate the burden of disease attributable to particulate matter ≤2.5 µm in aerodynamic diameter (PM_{2.5}). The IER function combines relative risk estimates from various PM25 sources, including active and passive smoking, to fill in the knowledge gap of air pollution studies in high exposure settings (Burnett et al. 2014). In the most recent GBD estimates (GBD 2019 Risk Factors Collaborators 2020), active smoking studies were excluded from the IER function to characterize risks at high exposure, because the few new studies of high air pollution conditions in Asia provided enough information so that evidence from active smoking data is no longer necessary to use. The number of studies of long-term air pollution and health in Asia,

however, remains limited to date, and there is a clear research gap with respect to the true size of the ambient air pollution and mortality associations in that region.

Dr. Vermeulen's study was funded through a special invitation based on several scientific and strategic considerations. At Utrecht University, the Netherlands, Dr. Vermeulen proposed to evaluate the association between long-term exposure to ambient air pollution and all-cause and cause-specific mortality in a pooled analysis of 23 Asian cohorts from the Asia Cohort Consortium (Aim 1). Moreover, he proposed to explore the heterogeneity in mortality risks among cohorts in the context of cultural, social, economic, or infrastructural differences between countries (Aim 2). Although the application came outside of a specific Request for Applications, it was reviewed using the same two-stage process: external reviewers evaluated the technical quality of the proposed work, followed by a discussion of strengths and limitations by the Research Committee. The HEI Research Committee recommended Dr. Vermeulen's application for funding because of the strong design features, the large number of participating cohorts, and the availability of individual-level covariate information. In addition, they appreciated that the cohorts were already harmonized, making it a cost-efficient and low-risk proposal. Dr. Vermeulen recruited Dr. George S. Downward as the analytical project lead.

During the course of the work, there were several unforeseen setbacks regarding cohort participation for various reasons, and only six of the original 23 cohorts that had expressed interest in participating were eventually included in the analyses. Therefore, the current report is focused solely on Aim 1. Aim 2 was not further pursued due to the small number of cohorts included in the final analyses.

This Commentary provides the HEI Review Committee's evaluation of the study. It is intended to aid the sponsors of HEI and the public by highlighting both the strengths and limitations of the study and by placing the Investigators' Report into a broader scientific perspective.

Dr. Roel Vermeulen's (principal investigator) 2-year study, "Long-Term Outdoor Air Pollution and Cause-Specific Mortality in a Pooled Analysis of 23 Asian Cohorts," began in July 2018. Total expenditures were \$236,000. The draft Investigators' Report from Downward (first author) and Vermeulen was received for review in September 2021. A revised report, received in August 2022, was accepted for publication in September 2022. During the review process, the HEI Review Committee and the investigators had the opportunity to exchange comments and to clarify issues in both the Investigators' Report and the Review Committee's Commentary.

This document has not been reviewed by public or private party institutions, including those that support the Health Effects Institute; therefore, it may not reflect the views of these parties, and no endorsements by them should be inferred.

^{*} A list of abbreviations and other terms appears at the end of this volume.

SCIENTIFIC BACKGROUND

The study by Downward and Vermeulen assessed the association between long-term exposure to ambient air pollution and all-cause and cause-specific mortality in an analysis of six Asian cohorts, with more than 340,000 participants (see Commentary Table 1 and Commentary Figure 1). The investigators estimated exposure to $PM_{2.5}$ and nitrogen dioxide (NO₂) at the residence of the participants for the year of recruitment using global satellite-based models. They applied single-pollutant Cox proportional hazard models to assess the association between air pollution exposure and all-cause and cause-specific mortality adjusted for important confounders, as described in more detail below.

STUDY POPULATION

The current study leverages the *Asia Cohort Consortium*, a multicenter collaborative effort since 2008 that consists of more than one million people to date from several dozen cohorts from 10 Asian countries. The Consortium seeks to understand the relation between genetics, environmental exposures, and the etiology of disease. To be eligible, cohorts must have information on mortality outcomes and important confounding variables, such as smoking and body mass index. Data on those variables were harmonized before entering the Consortium to ensure comparability.

The final analyses included six cohorts from the Asia Cohort Consortium and represented more than 340,000 adult participants in six countries (Commentary Figure 1, which includes the study name abbreviations). The study included three high-income countries (Japan, Taiwan, and Republic of Korea), one lower-middle country (Iran), and two low-income countries (Bangladesh, and India); designations are based on 2006 World Bank classifications. The cohorts were general population studies and varied widely in size, study period, recruitment method, geographical scope, exposure assignment, and outcome assessment (Commentary Table 1). The Indian MCS and the Japanese JPHC studies were the largest cohorts by far. Participants were recruited from 1991 to 2008 and followed-up between 5 and 23 years. Some cohorts were conducted in a single city or district (e.g., the Indian MCS and Bangladeshi HEALS), and others included much larger areas in a country (e.g., the Japanese JPHC). Mean exposures varied from 8 to 58 μ g/m³ for PM_{2.5} and 7 to 23 ppb for NO₂. Correlations between PM_{2.5} and NO₂ exposures varied from <0.01 to 0.57 (Commentary Table 1).

EXPOSURE ASSESSMENT

The investigators estimated exposure at the residence of the participants for $PM_{2.5}$ and NO_2 by using existing global satellite-based models and building on the exposure methods that were also used in the GBD project (Larkin et al. 2017; van Donkelaar et al. 2015, 2016). The global models provided high resolution (1 km² for $PM_{2.5}$, 100 m² for NO_2) annual average concentrations for 1998 (or 1997 in case of NO_2) to 2008. The

method is a sophisticated integration of primarily satellite data, with a chemical transport model, land-use information, and ground-monitoring data included as well. The models were validated against ground-based monitor data, with an overall R^2 of 0.81 and 0.54 for PM_{2.5} and NO₂, respectively. The estimates were assigned to study participants based on geocoded residential location data, but for the year of recruitment only. The 1998 exposure estimate was assigned for the participants that were recruited from 1991 to 1997 (i.e., before the global model estimates became available). Note that in four cohorts, exact address data were available for the year of recruitment; for the remaining cohorts (Indian MCS and Iranian Golestan) aggregated address data were used (e.g., postal codes).

OUTCOME ASSESSMENT

The study included both all-cause mortality and cause-specific mortality outcomes: nonaccidental, all cancer, lung cancer, cardiovascular disease, and noncancer lung disease mortality. The outcome assessment was performed by each individual cohort, typically through active follow-up or linkage to death registries. The same International Classification of Diseases (ICD) 9 or 10 coding was used for the different outcome categories across the cohorts except for the Japanese JPHC cohort. The JPHC cohort used slightly different ICD codes, particularly for cardiovascular disease. The JPHC cohort also did not have information on nonaccidental deaths.

ANALYSES

The investigators applied single-pollutant Cox proportional hazard models to assess the association between air pollution exposure and all-cause and cause-specific mortality. Models were adjusted for age (time axis), sex, recruitment year, smoking status, pack-years, body mass index, and a measure of socioeconomic status (education or employment). In addition, models adjusted for alcohol intake or diet for all cohorts except the Indian MCS and Bangladeshi HEALS cohorts. Models from one cohort (Iranian Golestan) were also adjusted for domestic fuel use — an indicator of household air pollution. That indicator was missing for the other cohorts.

The investigators calculated hazard ratios for each cohort separately and then combined using random effects meta-analysis. Associations were reported per 5- and 10-ppb increment in $PM_{2.5}$ and NO_2 , respectively. For each cohort, the investigators tested assumptions for the Cox proportional hazard models, ran two-pollutant models, and characterized the exposure–response function using splines and exposures by quartiles. Furthermore, they assessed the robustness of the associations by conducting several sensitivity and subgroup analyses. Notably, they conducted a sensitivity analysis in which associations were adjusted for urbanicity. Moreover, they reran analyses for the subcohorts of participants alive in 1998 when global model estimates became available. Note that no meta-analyses were conducted on any of the sensitivity analysis results.

Study Name	Location	Recruitment Years	Average Years of Follow-up	Sample Size	Mean Age	Mean Exposure PM _{2.5} (µg/m³)	Mean Exposure NO ₂ (ppb)	Correlation PM _{2.5} and NO ₂	Exposure Assignment	Outcome Assessment
MCS	Mumbai, India	1991– 1997	5	141,238	51	34	23	<0.01	Postal code	Active follow-up at regular intervals
JPHC	11 regions in Japan	1990– 1995	20	87,653	52	11	9	0.50	Residential address	Death registries
Golestan	Gonbad city and surround- ing rural area in Iran	2004– - 2008	11	49,982	52	32	9	0.54	Community level	Active follow-up at 1-year intervals
CBCSCP	7 townships in Taiwan	1991– 1992	23	23,759	47	8	9	0.14	Residential address	Health examina- tions, medical records, and cancer and death registries
HEALS	Araihazar in Bangladesh	2000– 2008	10	19,990	37	58	7	0.46	Residential address	Active follow-up at 1-year intervals
KMCC	4 areas in the Republic of Korea	1993– 2005	13	18,529	55	23	11	0.57	Residential address	Health insurance, cancer, and death registries
TOTAL	6 countries	1991– 2008	5-23	341,151	37-55	8-58	7–23	<0.01-0.57		

Commentary Table 1. Key Characteristics of the Six Asian Cohorts at Recruitment (ordered by sample size)



Commentary Figure 1. Geographical location of the six Asian cohorts.

SUMMARY OF RESULTS ACROSS COHORTS

- The meta-analytical summary effect estimates documented no association between long-term exposure to ambient PM_{2.5} and all-cause mortality and cause-specific mortality, except for a positive association with cardio-vascular mortality (Commentary Figure 2). The combined estimate for cardiovascular mortality was 1.05 per 5-µg/m³ increment and was borderline significant (95% confidence interval 0.99–1.12).
- For ambient NO₂, the combined estimates showed positive associations for all mortality outcomes, in particular the cancer outcomes. The combined estimate for all-cancer and lung cancer mortality were 1.18 and 1.13 per 10-ppb increment, respectively; both estimates were statistically significant. Combined estimates were heavily driven by positive associations from a single cohort (see below).

SUMMARY OF RESULTS WITHIN COHORTS

- The two largest cohorts the Indian MCS and the Japanese JPHC and the smaller Taiwanese CBCSCP cohort reported positive associations between ambient PM_{2.5} and cardiovascular mortality (Commentary Table 2). Those associations were statistically significant and fairly robust to further adjustment for urbanicity. The other three cohorts did not find an association with cardiovascular mortality.
- For ambient NO₂, the combined estimates for cancer outcomes were heavily influenced by the positive association in the Japanese JPHC cohort. This cohort carried

greater than 90% of the weight in meta-analyses. Most other cohorts documented no association with cancer outcomes.

• Large heterogeneity of the findings was reported across the cohorts, with null, negative, or positive findings, with sometimes no apparent pattern (Commentary Table 2). The Iranian Golestan and Korean KMCC cohorts consistently reported null findings. Findings from the Bangladeshi HEALS cohort were uninformative, due partly to the large confidence intervals and minimal exposure contrast. Hence, this cohort carried the lowest weight in the meta-analyses (often below 1%).

HEI REVIEW COMMITTEE'S EVALUATION

In its independent review of the study, the HEI Review Committee thought the research was well motivated and addressed a clear research gap. There are few long-term air pollution and health studies in Asia, and additional studies are urgently needed. This report adds to the overall knowledge base on health outcomes associated with air pollution in Asia. Although the number of cohorts participating was lower than anticipated when the study was funded, the inclusion of six harmonized cohorts ensured a large sample size.

In summary, the study documented large heterogeneity of the findings across the individual cohorts, with no association between long-term exposure to ambient $PM_{2.5}$ and all-cause mortality and cause-specific mortality in meta-analyses of all cohorts combined, except for a borderline significant positive association with cardiovascular mortality. Several individual cohorts (i.e., Indian MCS, Japanese JPHC, and Taiwanese





Commentary Figure 2. Meta-analysis for the association between exposure to PM_{2.5} (per 5 µg/m³) and NO₂ (per 10 ppb) and mortality in the six Asian cohorts combined. Associations were from single-pollutant models and adjusted for important potential confounders, such as smoking, body mass index, and socioeconomic status.

CBCSCP), however, did display positive significant associations between ambient $PM_{2.5}$ and cardiovascular mortality. For ambient NO_2 , the combined estimates showed positive associations for all mortality outcomes, in particular the cancer outcomes, although estimates were heavily driven by positive associations from the Japanese JPHC cohort.

The Committee noted several strengths of the research. First, it recognized the benefits of leveraging the Asia Cohort Consortium to study health effects of ambient air pollution. The study included data from six cohorts, representing more than 340,000 adult participants, which is a large sample size. The data were already harmonized and included both all-cause mortality and cause-specific mortality outcomes. There were also data available for several individual-level lifestyle factors, such as smoking status and intensity, body mass index, and diet, and the analyses were adjusted accordingly. As such, the study provides a useful model for future applications of harmonized cohort data to study the effects of air pollution on human health.

Second, the Committee appreciated the uniform assessment of long-term PM2.5 and NO2 using state-of-the-art exposure estimation methods. Exposures to PM2, and NO2 were estimated at a reasonably high spatial resolution - residential address level for most of the cohorts - and took advantage of global satellite-based models. The existing monitoring networks have limited spatial coverage with typically few stations in suburban and rural locations, particularly in lowand middle-income countries. According to the 2022 WHO Air Quality database, 40% of countries have no ground-level PM monitors. Ground-based monitor data are even sparser for NO₂, with 62% of countries with no monitors (WHO 2022). In addition, most existing monitoring networks have insufficient density to capture small-scale (within-city) variation of air pollution, which can be substantial for certain pollutants, such as NO_a.

Recent developments in satellite-based remote sensing and other exposure methods and models offer new ways to

			P	M _{2.5}		NO ₂						
	MCS	JPHC	Goles- tan	CBCSCP	HEALS	KMCC	MCS	JPHC	Goles- tan	CBCSCP	HEALS	KMCC
	India	Japan	Iran	Taiwan	Bangla- desh	Korea	India	Japan	Iran	Taiwan	Bangla- desh	Korea
All-cause	+	+	0	0	0	- (0)	+	+	0	_	+	0
Nonaccidental	+	NR	0	0 (+)	0	- (0)	+	NR	0	-	+	0
All-cancer	0	+	0	0 (+)	0	0	+	+	0	0	0	0
Lung cancer	0	0	0	0	0	0	0	+	0	0	0	0
Cardiovascular	+	+	0	+	0	0	+	+	0	0	+	0
Noncancer lung disease	0	-	0	0	0	0	0	0 (+)	0	- (0)	0	0

Commentary Table 2. Summary of Null, Negative, or Positive Findings in the Six Asian Cohorts

0 = null findings; - = negative association; + = positive association; NR = not reported. In parentheses, the summary of findings after additional adjustment for urbanicity when those findings differed from the main analysis. Note that in the Indian MCS cohort it was assumed that this cohort was fully urban, and no further adjustment was conducted.

provide air pollution estimates that cover large areas in a country, whole countries, or even multiple countries, with a sufficiently high degree of spatial resolution. The global satellite-based models applied in this study allowed exposure to be estimated for a large urban and rural population in six Asian countries. The Committee also thought the analyses were generally straightforward and clearly presented in the report. For example, the Committee appreciated the various sensitivity and subgroup analyses, including the additional adjustment for urbanicity in a sensitivity analysis.

Although the Review Committee broadly agreed with the investigators' conclusions, it identified limitations detailed below that should be considered when interpreting the results.

INADEQUATE ADJUSTMENT LIKELY FOR CHARACTERISTICS THAT CORRELATE WITH AIR POLLUTION AND MORTALITY

The Committee was concerned that residual confounding was likely in the main analyses due to inadequate adjustment for characteristics that correlate with air pollution and mortality, most notably socioeconomic status and urbanicity. These characteristics are likely related to both exposure and health, and difficult to fully capture based on the available indicators. Findings sometimes differed for models that adjusted for urbanicity as compared to those that did not (see Commentary Table 2). The Committee thought the authors should have adjusted for urbanicity in their main models instead of adjusting for urbanicity in a sensitivity analysis, even if there was some modest collinearity between air pollution estimates and urbanicity in some cohorts, as documented by the investigators. The Committee does appreciate the tables in the main text that compare the results with and without the urbanicity variable, additions made in response to earlier Committee comments.

The need for adjustment for urbanicity was also shown in the recent Prospective Urban and Rural Epidemiology (PURE) study (Hystad et al. 2020). The primary analyses adjusted for an indicator variable (urban or rural location). Models that further adjusted for "unmeasured differences between urban and rural areas within centers, as well as differences across centres" resulted in notable different results, especially for mortality. For example, the negative association between PM_{2 5} and all-cause mortality flipped to a positive association.

In the PURE study a notable negative association was also observed between markers of healthcare (hospital admissions or medication use) and deaths; this result suggests that poorer access to healthcare could be responsible, at least partly, for the higher mortality rates in low- and middle-income countries. Socioeconomic status and access to healthcare are closely related in many settings (Dagenais et al. 2020).

Since socioeconomic status influences where people live and is related to both exposure and health, this is often considered to be one of the most important confounders in air pollution epidemiology (Clark et al. 2014; Hajat et al. 2015; O'Neill et al. 2003). Additionally, there is evidence of differing correlations between socioeconomic status and air pollution exposure by location, highlighting the importance of adjusting for socioeconomic status based on the specific setting (Cesaroni et al. 2010; Hajat et al. 2013; Wang et al. 2022).

The current study adjusted for socioeconomic status in a fairly basic way with the use of an individual socioeconomic status indicator (i.e., education or employment) as a fixed covariate effect in the health model of the individual cohorts. The Committee thought that more effort to capture individual or area-level socioeconomic status in the study would have been beneficial.



HETEROGENEITY IN EFFECT ESTIMATES

The Committee noted that although the same exposure assessment and statistical methods were used, large heterogeneity of the findings was reported across the cohorts, with null, negative, or positive findings, with sometimes no apparent pattern. Some heterogeneity of the findings is expected, given the wide diversity of the six Asian cohorts. Heterogeneity is likely due, for example, to differences in populations, with different exposure levels, pollution sources and mixtures, time periods, age structure and follow-up times, socioeconomic status, urban-rural status, health status, access to healthcare, and outcome misclassification. Some specific differences across the cohorts were particularly striking, such as the low exposure contrast (Indian MCS and Bangladeshi HEALS), the low correlation between PM₂ and NO₂ (Indian MCS and Taiwan CBCSCP), the large percentage of illiterate population (Iranian Golestan), the short follow-up time (Indian MCS), the young study population (Bangladeshi HEALS), the rural location (Bangladeshi HEALS), particularly urban location (Indian MCS), and the low percentage of number of deaths, in particular for cancer (Indian MCS). Those and other differences could have contributed to the large heterogeneity of the findings observed in the current study.

In the systematic reviews underpinning the 2021 WHO Air Quality Guidelines for long-term exposure to $PM_{2.5}$ and NO_2 , a high degree of heterogeneity of the findings was also observed; this result was expected given that studies were included from across the globe (Chen and Hoek 2020; Huangfu and Atkinson 2020). Most of the heterogeneity in those studies, however, was due to heterogeneity in the magnitude of the positive association, not in the direction of the association (negative or positive). In particular, the negative associations in the current study are puzzling and run counter to the evidence base that documents clear evidence that long-term exposure to ambient air pollution is associated with increased mortality.

In the current study, a thorough evaluation of heterogeneity in mortality risks between cohorts in the context of cultural, social, economic, or infrastructural differences between countries was originally planned but was not pursued due to the small number of cohorts included in the final analyses. Although that decision is understandable given the data available to the investigators, the Committee would have been interested in better understanding potential sources of heterogeneity in the findings and noted that many questions have been unresolved.

Although the analyses were straightforward and clearly presented in the report, the study could have benefitted from a more detailed discussion and interpretation of all results, including the various sensitivity and subgroup analyses. For example, the added exposure–response function analysis was not tied together with the predetermined categorical analysis. Also, the Cox proportional hazards assumptions were violated for $PM_{2.5}$ (Indian MCS) and NO_2 (Japanese JPHC) for some mortality outcomes; the implications of which were not thor-

oughly addressed by the investigators. Also, an evaluation of potential selection bias due to the loss of several key cohorts from their original plans would have been useful. These and other issues limit what can be inferred from this study.

SUBSTANTIAL TEMPORAL AND SPATIAL MISALIGNMENT OF THE EXPOSURE DATA

The Committee had concerns about the exposure assessment approach because of the substantial temporal and spatial misalignment of the data. The study relies on an historical exposure assessment at recruitment that can be temporally misaligned with the health data by 5 to 23 years, depending on the cohort. Several issues of concern with the exposure assessment were noted by the Committee. First, the back extrapolations used for the participants before 1998 when the global model estimates became available to match the exact period of interest could introduce additional exposure error. The two largest cohorts (Indian MCS and Japanese JPHC) might be particularly affected by this misalignment, because those were also among the oldest cohorts. A sensitivity analysis in the subcohorts of participants alive in 1998 when global model estimates became available were generally consistent with the findings from the full cohorts, which was reassuring. Second, information on residential addresses after recruitment (i.e., moving history) was not available. Hence, residential mobility was not incorporated in the exposure assessment. Residential mobility can be substantial, especially in some low-and middle-income regions that are undergoing rapid urbanization in recent decades with population migration from rural to urban regions. Third, for a few cohorts (Indian MCS and Iranian Golestan) aggregated residential address data were used since individual address data were unavailable. That might be a particular issue for a pollutant such as NO₂, which is characterized by greater spatial variability than $PM_{2.5}$ and is influenced heavily by local emission sources. PM₂₅, in contrast, has long-range and secondary components and thus varies primarily at a regional level (Cyrys et al. 2012; Eeftens et al. 2012). More broadly, although the study applied state-of-the art exposure estimation methods with validated models, model performance differed regionally, with poorer PM_{2.5} performance in Asia compared to the global evaluation, as described by van Donkelaar and colleagues (2015, 2016). For NO₂, the model performance in Asia approximately matched the global evaluation estimate (Larkin et al. 2017). Nonetheless, in a later GBD application, NO₂ adjustments were made to correct the Larkin estimates for a "high bias in rural areas" (Anenberg et al. 2022). It should be noted that in model evaluations, estimates are compared to ground-based monitor data, but such evaluations are hampered by the paucity of ground-based monitors, with most of them located in urban areas of North America and Europe, as discussed in an earlier section. Although the Committee understands that Drs. Downward and Vermeulen made best use of the global exposure models available, the substantial temporal and spatial misalignment of the exposure data might have influenced the analysis of mortality outcomes in unpredictable ways.

HOUSEHOLD AIR POLLUTION WAS NOT EXAMINED

Like most other ambient air pollution and health studies, household air pollution was not examined in the current study. The Committee thought household air pollution might be a potential confounder or effect modifier. The investigators also alluded to that issue in the discussion of the findings. Household air pollution results from the burning of various fuels (coal, charcoal, wood, agricultural residue, animal dung, and kerosene, among others) for heating or for cooking using open fires or cookstoves with limited ventilation. Burning those fuels produces an array of pollutants that could harm human health, including PM_{2.5}, black carbon, and carbon monoxide. This practice is carried out by about half of the world's population, primarily from low- and middle-income countries. According to the most recent estimates from the GBD project, household air pollution contributes to about one third of the overall deaths linked to air pollution in 2019 (HEI 2020).

Only one cohort (Iranian Golestan) adjusted for domestic fuel use — an indicator of household air pollution. That indicator was missing for the other cohorts, unfortunately. The investigators reported consistent null findings between ambient air pollution and mortality for the Golestan cohort but found positive associations between some polluting fuel use (i.e., wood, kerosene, or "other" organic fuel) and mortality, that remained after adjusting for ambient PM_{2.5}. Similarly, in the PURE study, associations with solid fuel use for cooking and all-cause and cardiovascular mortality and morbidity were much more pronounced than the ambient PM_{2.5} associations (Hystad et al. 2019, 2020).

Investigating the complex interplay between household and ambient air pollution with health is difficult because household air pollution is typically not measured for large populations over long periods of time. Hence, most studies rely on use of fuel types as an indicator of household air pollution. The Committee welcomes the investigators' future work on this topic using the Asian Cohort Consortium as described by Hosgood and colleagues (2019).

BROADER CONTEXT OF AIR POLLUTION AND HEALTH IN ASIA

The current study adds to a small evidence base in Asia, where the levels of air pollution are often high, and the types and sources of air pollution markedly differ from those in high-income settings. Although cross-sectional or short-term health studies are increasingly available in Asia, there are few studies focused on long-term exposure to ambient air pollution (Baumgartner et al. 2020). The evidence base documenting clear evidence that long-term exposure to ambient air pollution is associated with increased mortality from all causes, cardiovascular disease, respiratory disease, and lung cancer continues to be dominated by studies from North America and Europe. The recent systematic reviews underpinning the 2021 WHO Air Quality Guidelines for $PM_{2.5}$ and NO_2 identified only a few long-term studies in Asia, and no single study from Africa, Central

America, or South America (Chen and Hoek 2020; Huangfu and Atkinson 2020). For example, only three studies from Asia entered the PM25 meta-analysis for all-cause and cardiovascular mortality (Tsjeng et al. 2015; Yang et al. 2018; Yin et al. 2017). Some studies of long-term exposure and morbidity and mortality in Asia emerged more recently (Commentary Table 3). Most of the studies from Asia documented a positive association between long-term exposure to PM25 and mortality outcomes, but there remains uncertainty about the true size of the PM₂₅ mortality relative risks. A recent study particularly relevant for the current study is the PURE study, which also used similar satellite-based global models (Hystad et al. 2020). The PURE study investigated the association between long-term exposure to PM_{2.5} and all-cause and cardiovascular mortality and morbidity in a large, pooled cohort of adults from 21 countries, with most of the study population residing in low- and middle-income countries. The PURE study reported that long-term exposure to PM25 was associated with increased risk for cardiovascular mortality and morbidity and adjusted for many important confounders, such as smoking, physical activity, socioeconomic status, urban or rural location and fuel use for cooking. No consistent association was observed for all-cause mortality and noncardiovascular mortality, and models were sensitive to adjustment for urbanicity, similar to the current study.

Given the paucity of studies in high air pollution settings, an IER function was developed for the GBD study to estimate mortality relative risks across the global exposure range for burden assessments. The function integrated four types of PM₂₅ exposures (outdoor PM₂₅, household air pollution, active smoking, and second-hand smoking) associated with cause-specific mortality (Burnett et al. 2014). In the most recent GBD estimates (GBD 2019 Risk Factors Collaborators 2020), active smoking studies were excluded from the IER function to characterize risks at high exposure, because the few new studies of high air pollution conditions in Asia provided enough information so that evidence from active smoking data is no longer necessary to use. This led to substantial increases in the relative risk curve for ischemic heart disease and stroke at the high end of the curve compared to the integrated curve that included active smoking studies. Notable increases in the relative risk curve were also reported in a PM₂ exposure–response function (global exposure mortality model [GEMM]), which was solely based on ambient $PM_{2.5}$ studies (Burnett et al. 2018). The use of GEMM resulted in burden estimates that were two to three times higher than those from the IER function. For the GEMM they included data from 41 cohorts in 16 different countries, including three studies from Asia (Tseng et al. 2015; Wong et al. 2015; Yin et al. 2017).

The differences in burden estimates reflect current uncertainty about key assumptions underlying the IER and GEMM models and therefore about the true size of the $PM_{2.5}$ mortality relative risks, particularly at the low- and high-end of the global exposure range (Burnett and Cohen 2020). The study by Downward and Vermeulen highlights the urgent need for future studies that could prove to be useful in reducing this

ΗE

Reference	Study Name	Location	Study Period	Sample Size	Mean PM _{2.5}	Mortality Outcome	Hazard Ratio per 5 µg/m³ª
Tseng et al. 2015	Civil servants' cohort	Greater Taipei, Taiwan	1989–2008	43,227	~29	All-cause Cardiovascular	0.96 (0.85–1.08) 0.89 (0.65–1.22)
Yin et al. 2017	Chinese men	45 districts in China	1990–2005	189,793	43.7	All-cause Cardiovascular Lung cancer	1.04 (1.04–1.05) 1.04 (1.04–1.05) 1.06 (1.04–1.08)
Yang et al. 2018	Hong Kong elderly	Hong Kong	1998–2011	66,820	42.2	All-cause Cardiovascular Respiratory	1.03 (1.01–1.05) 1.05 (1.02–1.09) 1.01 (0.97–1.06)
Li et al. 2018	CLHLS	China	2008–2014	13,344	50.7	All-cause	1.04 (1.03–1.05)
Yorifuji et al. 2019	Okayama City	Okayama City, Japan	2006–2016	75,569	14.0	All-cause Cardiovascular Lung cancer	1.29 (1.18–1.41) 1.06 (0.90–1.26) 1.63 (1.13–2.34)
Hystad et al. 2020	PURE	17 low- and middle-income countries	2003–2018	140,020	47.5 (all 21 countries)	All-cause Cardiovascular Noncardiovascular mortality Cardiovascular event (fatal + nonfatal)	0.99 (0.98–1.00) 1.02 (1.00–1.03) 0.98 (0.96–0.99) 1.03 (1.01–1.04)
Kim et al. 2020	NHIS-NSC	Republic of Korea	2002–2013	436,933	18.8	All-cause Cardiovascular	1.02 (1.01–1.02) 1.03 (1.02–1.03)
Brown et al. 2022	MDS	India	2004–2013	6.8 million	24.3	All-cause Ischemic heart disease Stroke Respiratory	1.01 (1.01–1.02) 1.00 (0.99–1.02) 1.04 (1.02–1.07) 1.01 (0.98–1.03)

Commentary Table 3. Summary of Selected Studies on Long-Term Exposure to PM_{2.5} and Mortality in Asia (in order of publication year)

CLHLS = Chinese Longitudinal Healthy Longevity Survey; MDS = Million Death Study; NHIS-NSC = National Health Insurance Service-National Sample Cohort; PURE = Prospective Urban and Rural Epidemiology (PURE).

^aFindings are converted to 5-µg/m³ increase in PM_{2.5} to allow comparison with the current study.

uncertainty. At some point in the near future with sufficient studies, it might be possible to develop separate risk curves for outdoor air pollution, second-hand smoking, and household air pollution in the GBD study. Having those separate risk curves would remove an important source of uncertainty related to equitoxicity of particles (assuming no differences in health impact by PM source, size, and chemical composition) as well as uncertainties related to some other aspects of exposure to those distinct sources of PM.

SUMMARY AND CONCLUSION

Drs. Downward and Vermeulen have assessed the association between long-term exposure to ambient air pollution and all-cause and cause-specific mortality in an analysis of six Asian cohorts. The research was well motivated and addressed a clear research gap. The large sample size and leverage of harmonized data from the Asia Cohort Consortium were considered to be strengths of the study. Furthermore, data were available for several individual-level lifestyle factors, such as smoking status and intensity, body mass index, and diet, and the analyses were adjusted accordingly. Application of existing global satellite-based models allowed for a uniform estimation of exposure at a reasonably high spatial resolution for a large urban and rural population in six Asian countries. Such a study would otherwise not have been possible given the paucity of ground-based monitors, particularly in low- and middle-income countries.

The study documented no association between longterm exposure to ambient PM25 and all-cause mortality and cause-specific mortality in meta-analyses, except for a borderline significant positive association with cardiovascular mortality. Several individual cohorts (i.e., Indian MCS, Japanese JPHC, and Taiwanese CBCSCP), however, did display positive significant associations between ambient PM,, and cardiovascular mortality. For ambient NO₂, the combined estimates showed positive associations for all mortality outcomes, in particular the cancer outcomes, although estimates were heavily driven by positive associations from the Japanese JPHC cohort. The cohorts were very diverse and large heterogeneity of the findings was reported across the individual cohorts, with null, negative, or positive findings, with sometimes no apparent pattern. Although the Review Committee broadly agreed with the investigators' conclusions, it identified limitations that should be considered when interpreting the results.

Importantly, the Committee was concerned that residual confounding was likely in the main analyses due to inadequate adjustment for characteristics that correlate with air pollution and mortality, most notably socioeconomic status and urbanicity. Findings sometimes differed for models that adjusted for urbanicity as compared to those that did not. The Committee would have been interested in better understanding potential sources of heterogeneity in the findings. There were also concerns about the exposure assessment approach because of the substantial temporal and spatial misalignment of the data, which might have influenced the analysis of mortality outcomes in unpredictable ways.

Overall, there remains uncertainty about the true size of the ambient air pollution and mortality associations in Asia, where the levels of air pollution are often high, and the types and sources of air pollution, including household air pollution, markedly differ from those in high-income settings. The study by Downward and Vermeulen highlights the urgent need for future studies that could prove to be useful in reducing this uncertainty. At the same time, these populations are experiencing very high levels of air pollution, meriting attention and action to reduce ambient air pollution regardless of the uncertainties.

ACKNOWLEDGMENTS

The Review Committee thanks the ad hoc reviewers for their help in evaluating the scientific merit of the Investigators' Report. The Committee is also grateful to Eleanne van Vliet and Pallavi Pant for oversight of the study; to Yi Lu for assistance with review of the draft report; to Hanna Boogaard for assistance with review of the report and in preparing its Commentary; to Mary Brennan for editing of this Report and its Commentary; and to Kristin Eckles and Hope Green for their roles in preparing this Research Report for publication.

REFERENCES

Anenberg SC, Mohegh A, Goldberg DL, Kerr GH, Brauer M, Burkart K, et al. 2022. Long-term trends in urban NO_2 concentrations and associated paediatric asthma incidence: Estimates from global datasets. Lancet Planet Health 6:e49–e58; doi:10.1016/S2542-5196(21)00255-2.

Baumgartner J, Brauer M, Ezzati M. 2020. The role of cities in reducing the cardiovascular impacts of environmental pollution in low- and middle-income countries. BMC Med 18:39; doi:10.1186/s12916-020-1499-y.

Brown P, Izawa Y, Balakrishnan K, Fu SH, Chakma J, Menon G, et al. 2022. Mortality from particulate matter 2.5 in India: National retrospective cohort study. Environ Health Perspect 130:097004; doi:10.1289/EHP10979.

Burnett R, Chen H, Szyszkowicz M, Fann N, Hubbell B, Pope CA 3rd, et al. 2018. Global estimates of mortality associated with long-term exposure to outdoor fine particulate matter. Proc Natl Acad Sci USA 115:9592–9597; doi:10.1073/pnas.1803222115.

Burnett R, Cohen A. 2020. Relative risk functions for estimating excess mortality attributable to outdoor $PM_{2.5}$ air pollution: Evolution and state-of-the-art. Atmosphere 11:589; doi:10.3390/atmos11060589.

Burnett RT, Pope CA 3rd, Ezzati M, Olives C, Lim SS, Mehta S, et al. 2014. An integrated risk function for estimating the global burden of disease attributable to ambient fine particulate matter exposure. Environ Health Perspect 122:397–403; doi:10.1289/ehp.1307049.

Cesaroni G, Badaloni C, Romano V, Donato E, Perucci CA, Forastiere F. 2010. Socioeconomic position and health status of people who live near busy roads: The Rome longitudinal study (RoLS). Environ Health 9:41; doi:10.1186/1476-069X-9-41.

Chen J, Hoek G. 2020. Long-term exposure to PM and allcause and cause-specific mortality: A systematic review and meta-analysis. Environ Int 143:105974; doi:10.1016/j. envint.2020.105974.

Clark LP, Millet DB, Marshall JD. 2014. National patterns in environmental injustice and inequality: Outdoor NO_2 air pollution in the United States. PLoS one 9:e94431; doi:10.1371/journal.pone.0094431.

Cyrys J, Eeftens M, Heinrich J, Ampe C, Armengaud A, Beelen R, et al. 2012. Variation of NO_2 and NO_x concentrations between and within 36 European study areas: Results from the ESCAPE study. Atmos Environ 62:374-390; doi:10.1016/j. atmosenv.2012.07.080.

Dagenais GR, Leong DP, Rangarajan S, Lanas F, Lopez-Jaramillo P, Gupta R, et al. 2020. Variations in common diseases, hospital admissions, and deaths in middle-aged adults in 21 countries from five continents (PURE): A prospective cohort study. Lancet 395:785–794; doi:10.1016/S0140-6736(19)32007-0.



Eeftens M, Tsai M-Y, Ampe C, Anwander B, Beelen R, Bellander T, et al. 2012. Spatial variation of $PM_{2.5}$, PM_{10} , $PM_{2.5}$ absorbance and PM-coarse concentrations between and within 20 European study areas and the relationship with NO₂: Results of the ESCAPE project. Atmos Environ 62:303–317; doi:10.1016/j.atmosenv.2012.08.038.

GBD 2019 Risk Factors Collaborators. 2020. Global burden of 87 risk factors in 204 countries and territories, 1990–2019: A systematic analysis for the Global Burden of Disease Study 2019. Lancet 396:1223–1249; doi:10.1016/S0140-6736(20)30752-2.

Hajat A, Diez-Roux AV, Adar SD, Auchincloss AH, Lovasi GS, O'Neill MS, et al. 2013. Air pollution and individual and neighborhood socioeconomic status: Evidence from the multi-ethnic study of atherosclerosis (MESA). Environ Health Perspect 121:1325–1333; doi:10.1289/ehp.1206337.

Hajat A, Hsia C, O'Neill MS. 2015. Socioeconomic disparities and air pollution exposure: A global review. Curr Environ Health Rep 2:440–450; doi:10.1007/s40572-015-0069-5.

HEI (Health Effects Institute). 2020. State of Global Air 2020. Special Report. Boston, MA:Health Effects Institute.

Hosgood HD III, Klugman M, Matsuo K, White AJ, Sadakane A, Shu XO et al. 2019. The establishment of the Household Air Pollution Consortium (HAPCO). Atmosphere (Basel) 10:10.3390/atmos10070422; doi:10.3390/atmos10070422.

Huangfu P, Atkinson R. 2020. Long-term exposure to NO_2 and O_3 and all-cause and respiratory mortality: A systematic review and meta-analysis. Environ Int 144:105998; doi:10.1016/j.envint.2020.105998.

Hystad P, Duong M, Brauer M, Larkin A, Arku R, Kurmi OP, et al. 2019. Health effects of household solid fuel use: findings from 11 countries within the Prospective Urban and Rural Epidemiology study. Environ Health Perspect 127:57003. doi:10.1289/EHP3915.

Hystad P, Larkin A, Rangarajan S, AlHabib KF, Avezum Á, Calik KBT, et al. 2020. Associations of outdoor fine particulate air pollution and cardiovascular disease in 157,436 individuals from 21 high-income, middle-income, and low-income countries (PURE): A prospective cohort study. Lancet Planet Health 4:e235–e245; doi:10.1016/S2542-5196(20)30103-0.

IARC (International Agency for Research on Cancer) Working Group on the Evaluation of Carcinogenic Risks to Humans. 2016. Outdoor air pollution. IARC Monogr Eval Carcinog Risks Hum 109:9–444.

Kim IS, Yang PS, Lee J, Yu HT, Kim TH, Uhm JS, et al. 2020. Long-term fine particulate matter exposure and cardiovascular mortality in the general population: A nationwide cohort study. J Cardiol 75:549–558; doi:10.1016/j.jjcc.2019.11.004.

Larkin A, Geddes JA, Martin RV, Xiao Q, Liu Y, Marshall JD, et al. 2017. Global land use regression model for nitrogen dioxide air pollution. Environ Sci Technol 51:6957–6964; doi:10.1021/acs.est.7b01148.

Li T, Zhang Y, Wang J, Xu D, Yin Z, Chen H, et al. 2018. Allcause mortality risk associated with long-term exposure to ambient $PM_{2.5}$ in China: A cohort study. Lancet Public Health 3:e470–e477; doi:10.1016/S2468-2667(18)30144-0.

O'Neill MS, Jerrett M, Kawachi L, Levy JL, Cohen AJ, Gouveia N, et al. 2003. Health, wealth, and air pollution: Advancing theory and methods. Environ Health Perspect 111:1861–1870; doi:10.1289/ehp.6334.

Thurston GD, Kipen H, Annesi-Maesano I, Balmes J, Brook RD, Cromar K, et al. 2017. A joint ERS/ATS policy statement: What constitutes an adverse health effect of air pollution? An analytical framework. Eur Respir J 49:1600419; doi:10.1183/13993003.00419-2016.

Tseng E, Ho WC, Lin MH, Cheng TJ, Chen PC, Lin HH. 2015. Chronic exposure to particulate matter and risk of cardiovascular mortality: Cohort study from Taiwan. BMC Public Health 15:936; doi:10.1186/s12889-015-2272-6.

U.S. EPA (U.S. Environmental Protection Agency). 2016. Integrated Science Assessment for Oxides of Nitrogen–Health Criteria. EPA/600/R-15/068. Washington, DC:U.S. EPA.

U.S. EPA (U.S. Environmental Protection Agency). 2019. Integrated Science Assessment for Particulate Matter (Final Report December 2019). EPA/600/R-19/188. Washington, DC:U.S. EPA.

van Donkelaar A, Martin RV, Brauer M, Boys BL. 2015. Use of satellite observations for long-term exposure assessment of global concentrations of fine particulate matter. Environ Health Perspect 1123:135–143; doi:10.1289/ehp.1408646

van Donkelaar A, Martin RV, Brauer M, Hsu NC, Kahn RA, Levy RC, et al. 2016. Global estimates of fine particulate matter using a combined geophysical-statistical method with information from satellites, models, and monitors. Environ Sci Technol 50:3762–3772; doi:10.1021/acs.est.5b05833.

Wang Y, Wang Y, Xu H, Zhao Y, Marshall JD. 2022. Ambient air pollution and socioeconomic status in China. Environ Health Perspect 130:67001; doi:10.1289/EHP9872.

WHO (World Health Organization). 2021. WHO Global Air Quality Guidelines: Particulate Matter ($PM_{2.5}$ and PM_{10}), Ozone, Nitrogen Dioxide, Sulfur Dioxide and Carbon Monoxide. WHO Regional Office for Europe. Geneva:World Health Organization.

WHO (World Health Organization). 2022. WHO Air Quality Database 2022. Available: https://www.who.int/publications/m/item/ who-air-quality-database-2022 [accessed January 24, 2023].

Wong CM, Lai HK, Tsang H, Thach TQ, Thomas GN, Lam KBH, et al. 2015. Satellite-based estimates of long-term exposure to fine particles and association with mortality in elderly Hong Kong residents. Environ Health Perspect 123:1167–1172; doi:10.1289/ehp.1408264.



Yang Y, Tang R, Qiu H, Lai PC, Wong P, Thach TQ, et al. 2018. Long-term exposure to air pollution and mortality in an elderly cohort in Hong Kong. Environ Int 117:99–106; doi:10.1016/j.envint.2018.04.034.

Yin P, Brauer M, Cohen A, Burnett RT, Liu J, Liu Y, et al. 2017. Long-term fine particulate matter exposure and nonaccidental and cause-specific mortality in a large national cohort of Chinese men. Environ Health Perspect 125:117002; doi:10.1289/ EHP1673.

Yorifuji T, Kashima S, Tani Y, Yamakawa J, Doi H. 2019. Longterm exposure to fine particulate matter and natural-cause and cause-specific mortality in Japan. Environ Epidemiol 3:e051; doi:10.1097/EE9.00000000000051.

ABBREVIATIONS AND OTHER ITEMS

AERONET	AErosol RObotic NETwork
ACC	Asia Cohort Consortium
CBCSCP	Community-based Cancer Screening Program
CI	confidence interval
DALYs	disability adjusted life years
ELAPSE	Effects of Low-Level Air Pollution: A Study in Europe
ESCAPE	European Study of Cohorts for Air Pollution Effects
GBD	Global Burden of Disease
GEMM	Global Exposure Mortality Model
HEALS	Health Effects for Arsenic Longitudinal Study
HIC	high-income country
HR	hazard ratio
ICD	International Classification of Diseases
IER	integrated exposure–response
JPHC	Japan Public Health Center-based Prospective Study
KMCC	Korean Multi-center Cancer Cohort Study
LMIC	low-and-middle-income countries
LUR	land use regression
MCS	Mumbai Cohort Study
NO_2	nitrogen dioxide
$PM_{2.5}$	particulate matter ≤2.5 µm in aerodynamic diameter
ppb	parts per billion
PURE	Prospective Urban and Rural Epidemiology
R^2	coefficient of determination
RR	relative risk
sd	standard deviation
WHO	World Health Organization

HEI BOARD, COMMITTEES, and STAFF

Board of Directors

Richard A. Meserve, Chair Senior of Counsel, Covington & Burling LLP; President Emeritus, Carnegie Institution for Science; former Chair, U.S. Nuclear Regulatory Commission

Jo Ivey Boufford Clinical Professor of Global Health and Pediatrics, New York University

Homer A. Boushey Emeritus Professor of Medicine, University of California–San Francisco

Jared L. Cohon President Emeritus and Professor, Civil and Environmental Engineering and Engineering and Public Policy, Carnegie Mellon University

Stephen Corman President, Corman Enterprises

Martha J. Crawford Operating Partner, Macquarie Asset Management

Ana V. Diez Roux Dana and David Dornsife Dean and Distinguished University Professor of Epidemiology, Dornsife School of Public Health, Drexel University; Director, Drexel Urban Health Collaborative

Michael J. Klag Dean Emeritus and Second Century Distinguished Professor, Johns Hopkins Bloomberg School of Public Health

Alan I. Leshner CEO Emeritus, American Association for the Advancement of Science

Catherine L. Ross Regents' Professor and Harry West Professor of City and Regional Planning and Civil and Environmental Engineering, and Director of the Center for Quality Growth and Regional Development, Georgia Institute of Technology

Martha E. Rudolph Environmental Attorney, Former Director of Environmental Programs, Colorado Department of Public Health and Environment

Karen C. Seto Frederick Hixon Professor of Geography and Urbanization Science, Yale School of the Environment

Research Committee

David A. Savitz, Chair Professor of Epidemiology, School of Public Health, and Professor of Obstetrics and Gynecology, Alpert Medical School, Brown University

David C. Dorman Professor, Department of Molecular Biomedical Sciences, College of Veterinary Medicine, North Carolina State University

Christina H. Fuller Associate Professor, School of Environmental, Civil, Agricultural and Mechanical Engineering, University of Georgia College of Engineering

Marianne Hatzopoulou Professor, Civil and Mineral Engineering, University of Toronto, Canada Research Chair in Transport Decarbonization and Air Quality

Amy H. Herring Sara & Charles Ayres Professor of Statistical Science and Global Health, Duke University

Heather A. Holmes Associate Professor, Department of Chemical Engineering, University of Utah

Neil Pearce Professor of Epidemiology and Biostatistics, London School of Hygiene and Tropical Medicine, United Kingdom

Ana M. Rule Assistant Professor and Director, Environmental Exposure Assessment Laboratories, Department of Environmental Health and Engineering, Johns Hopkins School of Public Health

Evangelia (Evi) Samoli Associate Professor of Epidemiology and Medical Statistics, Department of Hygiene, Epidemiology and Medical Statistics, School of Medicine, National and Kapodistrian University of Athens, Greece

Neeta Thakur Associate Professor of Medicine, University of California, San Francisco

Gregory Wellenius Professor, Department of Environmental Health, Boston University School of Public Health

Continues next page

HEI BOARD, COMMITTEES, and STAFF

Review Committee

Melissa J. Perry, Chair Dean, College of Health and Human Services, George Mason University

Sara D. Adar Associate Professor and Associate Chair, Department of Epidemiology, University of Michigan School of Public Health

Kiros T. Berhane Professor and Chair, Department of Biostatistics, Mailman School of Public Health, Columbia University

Ulrike Gehring Associate Professor, Institute for Risk Assessment Sciences, Utrecht University

Michael Jerrett Professor and Chair, Department of Environmental Health Sciences, Fielding School of Public Health, University of California–Los Angeles

Frank Kelly Henry Battcock Chair of Environment and Health and Director of the Environmental Research Group, Imperial College London School of Public Health, United Kingdom

Jana B. Milford Professor Emerita, Department of Mechanical Engineering and Environmental Engineering Program, University of Colorado–Boulder

Jennifer L. Peel Professor of Epidemiology, Colorado School of Public Health and Department of Environmental and Radiological Health Sciences, Colorado State University

Eric J. Tchetgen Tchetgen Luddy Family President's Distinguished Professor, Professor of Statistics and Data Science, The Wharton School, University of Pennsylvania

John Volckens Professor, Department of Mechanical Engineering, Walter Scott Jr. College of Engineering, Colorado State University

Officers and Staff

Daniel S. GreenbaumPresidentRobert M. O'KeefeVice PresidentEllen K. MantusDirector of ScienceDonna J. VorheesHEI Energy CEO and Vice PresidentAnnemoon M. van ErpDeputy Director of ScienceThomas J. ChampouxDirector of Science CommunicationsJacqueline C. RutledgeDirector of Finance and AdministrationJason DesmondDeputy Director of Finance and AdministrationEmily AldenCorporate Secretary

Amy AndreiniScience Communications SpecialistAyusha ArianaResearch AssistantPalak BalyanConsulting Staff ScientistHanna BoogaardConsulting Principal ScientistAaron J. CohenConsulting Principal ScientistDan CrouseSenior ScientistRobert M. DavidsonStaff AccountantPhilip J. DeMarcoCompliance Manager

Continues next page

HEI BOARD, COMMITTEES, and STAFF

Kristin C. Eckles Senior Editorial Manager Elise G. Elliott Staff Scientist Hope Green Editorial Project Manager Lissa McBurney Senior Science Administrator Janet I. McGovern Executive Assistant Victor Nthusi Consulting Research Fellow Pallavi Pant Head of Global Health Allison P. Patton Senior Scientist Quoc Pham Science Administrative Assistant Anna S. Rosofsky Senior Scientist Robert A. Shavers Operations Manager Eva Tanner Staff Scientist Alexis Vaskas Digital Communications Manager Ada Wright Research Assistant



H E A L T H E F F E C T S INSTITUTE

75 Federal Street, Suite 1400 Boston, MA 02110, USA +1-617-488-2300 www.healtheffects.org

RESEARCH REPORT

Number 213 May 2023 Publishing history: This document was posted at www.healtheffects.org in May 2023.

Citation for document:

Downward GS, Vermeulen R; Asia Cohort Consortium Executive Board. 2023. Ambient Air Pollution and All-Cause and Cause-Specific Mortality in an Analysis of Asian Cohorts. Research Report 213. Boston, MA:Health Effects Institute.

© 2023 Health Effects Institute, Boston, Mass., U.S.A. David Wade, Virginia Beach, Va., Compositor. Library of Congress Catalog Number for the HEI Report Series: WA 754 R432.